

# Falls-associated elevation of troponin level in an older woman with normal coronary arteries

## CASE REPORT

JKH Luk MBBS(HK), MSc (Experimental Medicine) (UBC), FRCP (Edin, Glasg, Ire), FHKCP, FHKAM(Medicine)

### ABSTRACT

A 75-year old woman with no history of ischaemic heart disease sustained a fall with scalp laceration. She had no chest pain. Troponin I was found to be high on admission and remained so for one day. Computed tomographic coronary angiogram was normal. She was discharged on low dose aspirin. The issue of non-ischaemic elevation of troponin is discussed.

*Correspondence to: Dr James KH Luk, Department of Medicine and Geriatrics, Fung Yiu King Hospital, 9 Sandy Bay Road, Pokfulam, Hong Kong. Email: lukkh@ha.org.hk*

### INTRODUCTION

Troponins are highly sensitive biomarkers of myocardial injury. They are useful in the diagnosis of acute myocardial infarction and risk stratification of acute coronary syndrome.<sup>1</sup> Nonetheless, troponin level may be raised in many conditions other than acute myocardial infarction, posing a diagnostic difficulty for clinicians.<sup>2,3</sup> We report an older woman who fell and sustained a scalp laceration. Troponin I was found to be high on admission; further investigation revealed no coronary artery disease.

### CASE REPORT

In March 2015, a 75-year old woman was taken by ambulance to the emergency department with a head injury and scalp laceration secondary to a non-syncopal fall. She had normal cognitive function and no history of ischaemic heart disease, diabetes mellitus, or hypertension. She was a non-smoker and non-drinker, and was independent in walking and activities of daily living. She had been taking simvastatin 10 mg nocte for hyperlipidaemia for 3 years. The patient reported the fall to be accidental when she missed the curb of the street. She sustained a 5 cm long laceration on her left forehead. She had no loss of consciousness or any other symptoms or injury.

Computed tomography (CT) of the brain revealed

no intracranial haemorrhage or skull fracture. In the ward, she was noted to be very nervous with a blood pressure of 200/100 mm Hg and a heart rate of 100 beats per minute. After a period of rest, her blood pressure settled to around 140/70 mm Hg and her heart rate dropped to 80 beats per minute.

Blood tests on admission revealed a raised troponin I level of 0.83 ng/ml (cut-off value for acute myocardial infarction, 0.5 ng/ml with 96% sensitivity and 94% specificity), and creatine phosphokinase (CPK) of 145 U/L (normal range, 40-160 U/L). Her renal and liver function tests were normal with a creatinine level of 53 umol/L. Electrocardiography demonstrated sinus rhythm and no Q wave or any ST or T changes. Because of the raised troponin I, she was started on low molecular weight heparin and aspirin 80 mg daily.

Subsequent blood tests showed that the troponin I level peaked at 2.22 ng/ml 6 hours after the first test and CPK remained static at 120 U/L. The troponin I level subsequently dropped to 1.9 ng/ml at 12 hours and 0.6 ng/ml at 18 hours.

She was seen by a cardiologist, and echocardiography showed normal contractility with ejection fraction of 60% and no valvular lesion. To exclude coronary heart disease, CT coronary angiogram was performed on day 2 and showed completely normal coronary arteries with an Agatston

score of 0. This suggests an extremely low likelihood of any coronary artery disease in an asymptomatic individual.<sup>4</sup> Low molecular weight heparin was stopped after 3 days, but aspirin 80 mg daily was continued. She was then seen by a geriatrician who carried out a comprehensive geriatric assessment. She was advised to perform balance exercises such as Tai Chi and was scheduled for follow-up in 12 weeks. The patient was discharged on day 3 and remained well.

## DISCUSSION

Falls and imbalance occur commonly in older people and are a major topic in geriatric medicine.<sup>5,6</sup> In Hong Kong, the prevalence of older people having at least one fall in the preceding 12 months ranges from 18% to 19.3%, with 75.2% sustaining an injury and 7.2% a serious injury.<sup>7,8</sup> Non-ischaemic cause of troponin elevation after falls has not been reported.

Troponins are regulatory proteins that control the calcium-mediated interaction of actin and myosin. The troponin complex consists of 3 subunits: troponin T, troponin I, and troponin C.<sup>9</sup> The skeletal and cardiac isoforms of troponin T and troponin I, but not troponin C, are distinct, and skeletal isoforms are not detected by the monoclonal antibody-based assays currently in use. Although troponin is sensitive and specific for detection of myocardial injury, it can be elevated in other conditions and is not specific for myocardial infarction.<sup>10</sup> Non-ischaemic causes of troponin elevation include sepsis, pulmonary embolism, renal failure, heart failure, pericarditis, cerebrovascular accident, tachycardia, Takotsubo cardiomyopathy, endurance exercise, angioplasty, cardiac contusion, and chemotherapy. The exact mechanism of non-ischaemic troponin elevation is complex and not well known. Troponins have a cytoplasmic pool and structural pool.<sup>11</sup> Myocardial injury first leads to release of the cytoplasmic pool resulting in a peak serum concentration at 24 to 36 hours after injury. The half-life of troponin in serum is around 2 hours, and the continuous elevation of troponin for a week or more depends on gradual degradation of the myofibrils and release of troponin from the structural pool as a result of myocardial injury. The mechanism for non-ischaemic troponin elevation is heterogeneous and multifactorial. In tachycardia, it is thought to be the result of shortening of diastole with sub-endocardial ischaemia.<sup>12</sup>

Troponin raised after angioplasty might be the result of procedural complications such as side branch occlusion and thrombus formation.<sup>13</sup> In heart failure, apoptosis, myocyte damage from neurohumoral activation and inflammatory cytokines might lead to troponin increase.<sup>14</sup> Acute right ventricular strain secondary to increased pulmonary artery stenosis, and hypoxaemia and hypoperfusion secondary to low cardiac output are postulated causes of raised troponin.<sup>15</sup> In Takotsubo cardiomyopathy, physical or emotional stress leads to increased circulatory catecholamines and myocardial stunning and surge in troponin.<sup>16</sup> In pericarditis, there might be inflammatory involvement of the epicardium with consequent myocyte damage.<sup>17</sup> Inflammatory cytokines, hypotension, tachycardia, and increased catecholamines can act together and result in raised blood levels of troponin. After endurance exercise, a rise in troponin level may be due to the release of cytoplasmic troponin or increased permeability of myocyte cells under stress.<sup>18</sup> The reason for a rise in troponin in renal failure remains unknown.<sup>19</sup>

The exact mechanism of raised troponin in this patient was not known. There was no evidence of coronary artery disease on CT coronary angiogram, and her raised troponin subsided quickly in less than one day, probably owing to the release of cytoplasmic troponin with change of myocyte permeability as a result of catecholamine surge. She was quite nervous on the day of admission with high blood pressure and tachycardia but these subsided after resting. Probably because of the lack of myocardial cell injury, troponin I level dropped within a day. Other possible mechanisms of a transient rise in troponin include an imbalance of oxygen supply and demand secondary to sudden hypertension and tachycardia, shortening of diastole with sub-endocardial ischaemia associated with tachycardia, small vessel disease of the heart, and myocardial ischaemia secondary to coronary spasm. None of these conditions would be excluded by a normal CT coronary angiogram.

CPK is an enzyme found mainly in the heart, brain, and skeletal muscle. In the blood, CPK may be an indication of damage to CPK-rich tissue, such as in rhabdomyolysis, myocardial infarction, myositis, and myocarditis. In our patient, CPK was not elevated despite the raised troponin I level, indicating the absence of significant myocyte damage.

There are currently no data from studies evaluating the therapies aimed at reducing risk in patients with non-ischaemic troponin elevation. In most cases, aggressive antithrombotic and antiplatelet treatments are not indicated. Aspirin, unless it is contraindicated, may be given. Whether to check the troponin level in this patient was the clinical judgment of the admitting doctor. Atypical presentation of an older patient is common and well reported, often requiring comprehensive geriatric assessment supplemented by appropriate investigations to delineate the underlying problems.<sup>20,21</sup> Although the patient did not complain of any chest discomfort on admission, if the admitting doctor was concerned that a cardiac event might have occurred leading to the 'trip and fall', or that a heart event might have occurred after the fall, checking the troponin I level would be helpful to exclude such coronary event. Progressive troponin I rise in this patient with cardiovascular risk factors (age and hyperlipidaemia) mandated further tests such as CT coronary angiogram to exclude significant coronary artery disease. Although CT coronary angiogram has a small risk of contrast nephropathy and allergy, it is less invasive than conventional cardiac catheterisation.

## CONCLUSION

Troponin may be elevated in the absence of acute myocardial infarction. Many conditions are associated with non-ischaemic troponin elevation, and their mechanisms are heterogeneous. There are currently no data from studies evaluating the therapies aimed at reducing risk in patients with non-ischaemic troponin elevation. Aspirin, unless contraindicated, can be given, as it appears to be relatively safe in most clinical circumstances.

## REFERENCES

- Rosenbaum LS, Januzzi JL. Moving troponin testing into the 21st century: will greater sensitivity be met with greater sensibility? *Cardiovasc Hematol Disord Drug Targets* 2008;8:118-26.
- Rahman A, Broadley SA. Review article: elevated troponin: diagnostic gold or fool's gold? *Emerg Med Australas* 2014;26:125-30.
- Mahajan N, Mehta Y, Rose M, Shani J, Lichstein E. Elevated troponin level is not synonymous with myocardial infarction. *Int J Cardiol* 2006;111:442-9.
- Abunassar JG, Yam Y, Chen L, D'Mello N, Chow BJ. Usefulness of the Agatston score = 0 to exclude ischemic cardiomyopathy in patients with heart failure. *Am J Cardiol* 2011;107:428-32.
- Luk JK, Chan TY, Chan DK. Fall prevention in the elderly: translating evidence into practice. *Hong Kong Med J* 2015;21:165-71.
- Willeboordse F, Hugtenburg JG, van Dijk L, et al. Opti-Med: the effectiveness of optimised clinical medication reviews in older people with 'geriatric giants' in general practice; study protocol of a cluster randomised controlled trial. *BMC Geriatr* 2014;14:116.
- Chu LW, Chi I, Chiu AY. Falls and fall-related injuries in community-dwelling elderly persons in Hong Kong: a study on risk factors, functional decline, and health services utilization after falls. *Hong Kong Med J* 2007;13(Suppl 1):S8-12.
- Chu LW, Chi I, Chiu AY. Incidence and predictors of falls in the Chinese elderly. *Ann Acad Med Singapore* 2005;34:60-72.
- Sevrieva I, Knowles AC, Kampourakis T, Sun YB. Regulatory domain of troponin moves dynamically during activation of cardiac muscle. *J Mol Cell Cardiol* 2014;75:181-7.
- Jeremias A, Gibson CM. Narrative review: alternative causes for elevated cardiac troponin levels when acute coronary syndromes are excluded. *Ann Intern Med* 2005;142:786-91.
- Remppis A, Scheffold T, Greten J, et al. Intracellular compartmentation of troponin T: release kinetics after global ischemia and calcium paradox in the isolated perfused rat heart. *J Mol Cell Cardiol* 1995;27:793-803.
- Ben Yedder N, Roux JF, Paredes FA. Troponin elevation in supraventricular tachycardia: primary dependence on heart rate. *Can J Cardiol* 2011;27:105-9.
- Prasad A, Rihal CS, Lennon RJ, Singh M, Jaffe AS, Holmes DR Jr. Significance of periprocedural myonecrosis on outcomes after percutaneous coronary intervention: an analysis of preintervention and postintervention troponin T levels in 5487 patients. *Circ Cardiovasc Interv* 2008;1:10-9.
- Felker GM, Mentz RJ, Teerlink JR, et al. Serial high sensitivity cardiac troponin T measurement in acute heart failure: insights from the RELAX-AHF study. *Eur J Heart Fail* 2015;17:1262-70.
- Ng PL, Kwok CH, Tsui SH, Tong HK. Raised troponin I in Chinese patients with acute pulmonary embolism suggests complicated clinical course and may predict intensive care unit admission. *Singapore Med J* 2013;54:86-9.
- Abdulla I, Ward MR. Tako-tsubo cardiomyopathy: how stress can mimic acute coronary occlusion. *Med J Aust* 2007;187:357-60.
- Newby LK, Ohman EM. Troponins in pericarditis: implications for diagnosis and management of chest pain patients. *Eur Heart J* 2000;21:798-800.
- Wedin JO, Henriksson AE. Postgame elevation of cardiac markers among elite floorball players. *Scand J Med Sci Sports* 2015;25:495-500.
- Mbagaya W, Luvai A, Lopez B. Biological variation of cardiac troponin in stable haemodialysis patients. *Ann Clin Biochem* 2015;52:562-8.
- Luk JK, Or KH, Woo J. Using the comprehensive geriatric assessment technique to assess elderly patients. *Hong Kong Med J* 2000;6:93-8.
- Limpawattana P, Phungoen P, Mitsungnern T, Laosuangkoon W, Tansangworn N. Atypical presentations of older adults at the emergency department and associated factors. *Arch Gerontol Geriatr* 2016;62:97-102.

OUTSTANDING FREE PAPERS

**Communication with older Chinese patients on advanced directives: is it feasible?**

Chan F, Chiu KC, Kwan SK, Chu LW, Chan HW

Acute Geriatric Unit, Grantham Hospital, Hong Kong West Cluster

**Background:** Advance directive (AD) is based on the principle of patients' autonomy in healthcare decisions. In Chinese culture, death is a sensitive issue, and overt reference to death can be regarded as disrespectful. Local study showed general acceptance of the AD in patients with advance cancers. Nonetheless, little is known in elderly patients without malignancy in this aspect. **Objective:** This study examined the acceptance rate and predictive factors of accepting the AD through communication of real-life experience with older patients about AD. **Methods:** We recruited 60 medically and mentally stable Chinese patients over 65 years of age from the Acute Geriatric Unit of the Grantham Hospital. The geriatricians educated and interviewed the patient and their relatives, and recorded the patient's preferences, and the refusal of futile treatment in case they suffered from terminal disease, persistent vegetative state, or irreversible coma, using an AD document designed by Hospital Authority. We documented baseline socio-demographic, medical, functional, and cognitive factors, and the reasons for accepting or rejecting the AD. **Results:** Amongst the 60 patients, 35% accepted and 65% rejected the AD. Factors that predicted AD acceptance included higher education level ( $p=0.046$ ), single or widow marital status ( $p<0.001$ ), and poor health status ( $p=0.028$ ). Factors that predicted AD rejection included independent functioning ( $p=0.039$ ), having children ( $p=0.007$ ), good social support ( $p=0.001$ ), and having family as the main carer ( $p=0.008$ ). The commonest reasons for AD acceptance were to avoid suffering (67%) and uphold quality of life above longevity (33%); and for rejecting AD were the belief that their family would make the decision (67%) and nature would take its course (31%). 23% remained unfamiliar with the concept of AD and 23% were not ready to discuss or accept it. **Discussion:** Only 35% of our patients engaged an AD, which is lower than the rates of 41% to 63% reported in studies on advance cancer patients. This may be because our subjects had better prognosis than cancer patients overall. Patients with poor social support (lives alone, widow or no children) and self-perception of poor health status were more likely to engage in AD, this may be related to above stressors alert the seniors to think more about end-of-life issue. Patients with good family support (patients with children or spouse/children as main carer) and good functional status were less likely to sign an AD. They believed that family members would decide for their

future care plan in case they were severely ill or mentally incompetent. These findings are consistent with traditional Chinese culture that children play an important role in older patient's treatment plan. **Conclusion:** AD is feasible in local older Chinese patients, but the engagement rate is not high. More public education on end-of-life issues is needed before AD can be widely adopted.

**Risk factors for prolonged dysphagia following acute stroke**

Lau WS

Department of Medicine and Geriatrics, Princess Margaret Hospital

**Background:** Dysphagia is a serious complication of stroke and is closely linked with aspiration and pneumonia. Factors influencing its recovery are not well known. This study aimed to determine the prevalence of post-stroke dysphagia, identify possible predictive factors and prognostic variables affecting swallowing recovery, and investigate its relationship with pneumonia and mortality 6 months after stroke. **Methods:** We retrospectively reviewed 190 patients admitted to rehabilitation ward with acute stroke between June 2013 and December 2013. Clinical findings, investigation results, and diet composition were collected. Subjects were divided into 3 groups: non-dysphagia ( $n=47$ ); transient dysphagia, diet normalised upon discharge ( $n=103$ ); and prolonged dysphagia, required modified diet upon discharge ( $n=40$ ). Welch's one-way analysis of variance, Chi-square analysis, and Holm-Bonferroni method were used. **Results:** Glasgow Coma Scale (GCS), modified Barthel Index (BI), Modified Rivermead Mobility Index (MRMI), and Modified Functional Ambulation Classification (MFAC) were significantly lower in the prolonged dysphagia group than in the transient dysphagia group ( $p<0.001$ ). Although hemispatial neglect was an independent risk factor for prolonged dysphagia ( $p=0.0186$ ), lesion location including left hemispheric, right hemispheric, bihemispheric disease, and aetiology of stroke did not significantly differ between groups. Patients with prolonged dysphagia had a 6-month mortality rate of 17.5%, and 42.5% of them suffered from all-cause pneumonia within 6 months. Hospital stay was significantly longer in transient and prolonged dysphagia patients ( $p<0.001$ ). **Conclusion:** Prolonged dysphagia was associated with lower initial body weight ( $p=0.01$ ) and increased stroke severity (evidenced by an initial lower GCS, modified BI, MRMI, MFAC, National Institutes of Health Stroke Scale, facial weakness, dysarthria, and hemispatial neglect). Patients with prolonged dysphagia were predicted to have a longer hospital stay, higher incidence of pneumonia and mortality, and a higher proportion moving to aged home upon discharge ( $p<0.001$ ).